

TOP MILK DAIRY 17

CLASS: Concentrated Dairy Feed 17

REG. NO. ACT 36/1947: V30874

Ideal supplement to sustain production of medium producing dairy cows on a high roughage diet or TMR. Perfectly balanced to supply the full spectrum of nutrients with 17% protein.



COMPOSITION

NUTRIENT	MIN/MAX	QUANTITY	UNIT
Protein (Of which a maximum of 30% originates from NPN)	Min	170	g/kg
Moisture	Max	120	g/kg
Fibre	Max	120	g/kg
Fat	Min/Max	25/70	g/kg
Calcium	Min/Max	10 /12	g/kg
Phosphorus	Min	5	g/kg
Urea	Max	14.81	g/kg

INGREDIENT STATEMENT:

This animal feed contains: Grain and grain by-products, animal protein products, plant protein products, roughages and agro-industrial by-products, sugar cane by-products, oils and fats, amino acids, vitamins and minerals, NPN Sources, antioxidants, binders, acidity regulators, flavouring agents, aroma enhancing agents, compounds of trace elements, digestibility enhancers and gut flora stabilizers. This product contains genetically modified ingredients.

NOTE:

This farm feed is available with or without registered stock remedies. See bag tag for details and feed strictly according to given instructions.

UREA CAUTION:

This farm feed contains feed grade UREA and/or other NPN sources and must therefore be fed strictly according to the instructions for use.

NPN WARNING:

1. Vinegar is an effective remedy against NPN poisoning. Mix with an equal amount of water. Dose half a bottle per calf or large sheep or 2 – 4 bottles per head of cattle. (1 bottle = 750ml).
2. Protect this farm feed against rain. NPN is soluble and animals drinking such a solution could be poisoned.
3. Do not feed this farm feed indiscriminately with other NPN containing farm feed. Consult an animal scientist.
4. Adaption: When changing from a NPN-free meal to a NPN-containing meal it is advisable to feed a 50/50 mixture over a period of 4 – 6 days.
5. Dairy feed must be fed 2 or more times daily.

FEEDING RECOMMENDATIONS:

Feed with roughage or grazing, quantities depending on:

1. Quality and availability of grazing and/or roughage;
2. Stage of lactation
3. Milk and butterfat production

General recommendation 500 gram/litre milk.



For more detailed feeding recommendations contact an Epol Technical Advisor.

